



Child's name _____ Birthdate _____

		Birth	1 month	2 months	4 months	6 months	9 months	12 months	15 months	18 months	19-23 months	2-3 years	4-6 years	7-10 years	11-12 years		
Recommended Immunizations	Disease	Vaccination	Visit date: _____	Visit date: _____	Visit date: _____	Visit date: _____	Visit date: _____	Visit date: _____	Visit date: _____	Visit date: _____	Visit date: _____	Visit date: _____	Visit date: _____	Visit date: _____	Visit date: _____		
	Hepatitis B	HepB	<input type="checkbox"/> 1st dose	<input type="checkbox"/> 2nd dose		<input type="checkbox"/> 3rd dose											
	Rotavirus	RV ¹		<input type="checkbox"/> 1st dose	<input type="checkbox"/> 2nd dose	See footnote ¹											
	Diphtheria, tetanus, whooping cough (pertussis)	DTaP – for children under age 7		<input type="checkbox"/> 1st dose	<input type="checkbox"/> 2nd dose	<input type="checkbox"/> 3rd dose		<input type="checkbox"/> 4th dose				<input type="checkbox"/> 5th dose					
	Haemophilus influenzae type b (bacterial meningitis)	Hib ²		<input type="checkbox"/> 1st dose	<input type="checkbox"/> 2nd dose	See footnote ²		3rd or 4th dose, see footnote ²									
	Pneumococcal	PCV13		<input type="checkbox"/> 1st dose	<input type="checkbox"/> 2nd dose	<input type="checkbox"/> 3rd dose		<input type="checkbox"/> 4th dose									
	Polio	IPV, inactivated		<input type="checkbox"/> 1st dose	<input type="checkbox"/> 2nd dose	<input type="checkbox"/> 3rd dose								<input type="checkbox"/> 4th dose			
	Flu (influenza)	Flu ³					<input type="checkbox"/> Vaccination every annual flu season, one or two doses, see footnote ³										
	Measles, mumps, rubella	MMR						<input type="checkbox"/> 1st dose						<input type="checkbox"/> 2nd dose			
	Chickenpox (varicella)	VAR						<input type="checkbox"/> 1st dose						<input type="checkbox"/> 2nd dose			
	Hepatitis A	HepA ⁴						Two-dose series, see footnote ⁴									
Meningococcal	Meningococcal																
Tetanus, diphtheria, whooping cough (pertussis)	Tdap – for children 7 years or older														1st dose, see footnote ⁶		
HPV (human papillomavirus)	HPV ⁵														See footnote ⁵		
Growth	At each well-child visit, enter date, length, weight and percentile information to keep track of your child's progress.		Weight/Percentile	Weight/Percentile	Weight/Percentile	Weight/Percentile	Weight/Percentile	Weight/Percentile	Weight/Percentile	Weight/Percentile	Weight/Percentile	Weight/Percentile	Weight/Percentile	Weight	Weight	Weight	
			Length/Percentile	Length/Percentile	Length/Percentile	Length/Percentile	Length/Percentile	Length/Percentile	Length/Percentile	Length/Percentile	Length/Percentile	Length/Percentile	Length/Percentile	Length/Percentile	Height	Height	Height
			Head circumference	Head circumference	Head circumference	Head circumference	Head circumference	Head circumference	Head circumference	Head circumference	Head circumference	Head circumference	Head circumference	Head circumference	BMI percentile	BMI percentile	BMI percentile

Shaded box indicates the vaccine can be given during shown age range.

If your child misses a shot, you don't need to start over. Just go back to your child's doctor for the next shot. If you have questions about vaccines, talk with your child's doctor.

¹RV: If RotarixR is used, a child should get a two-dose series at ages 2 and 4 months. If RotaTeqR is used, it should be a three-dose series at ages 2, 4 and 6 months. If any dose in the series was RotaTeq or if the vaccine product is not known for any dose, give a total of three doses of the rotavirus vaccine.

²Hib: Children will get a primary series of two or three doses and a booster dose. The booster will be dose three or four, depending on the vaccine used in the primary series.

³Flu: Children ages 6 months through 8 years getting the flu vaccine for the first time should get two doses at least four weeks apart.

⁴HepA: Two doses of the HepA vaccine are needed for lasting protection. The first dose should be given between 12 months and 23 months of age. The second dose should be given six to 18 months later.

⁵HPV: The vaccination series can start at age 9. If it is started before age 15, two doses are given six to 12 months apart. If immunization starts after age 15, three doses are recommended.

⁶Meningococcal: This is a 2-dose series. The second dose is given at age 16.