




# Keep Track of Your Baby's VACCINATIONS

Immunizations are the best way to make sure your child is protected against contagious diseases. Bring this to your child's appointments to track immunizations and milestones.



Child's name \_\_\_\_\_  
 Birthdate \_\_\_\_\_

## RECOMMENDED IMMUNIZATIONS

	Birth	1 month	2 months	4 months	6 months	9 months	12 months	15 months	18 months	23 months
<b>Hepatitis B (HepB)</b> <i>Protects against hepatitis B</i>	<input type="checkbox"/> 1st dose	<input type="checkbox"/> 2nd dose			<input type="checkbox"/> 3rd dose					
<b>Rotavirus<sup>1</sup> (RV)</b> <i>Protects against rotavirus</i>			<input type="checkbox"/> 1st dose	<input type="checkbox"/> 2nd dose	<input type="checkbox"/> See footnote <sup>1</sup>					
<b>Diphtheria, tetanus, pertussis (DTaP)</b> <i>Protects against diphtheria, tetanus and whooping cough</i>			<input type="checkbox"/> 1st dose	<input type="checkbox"/> 2nd dose	<input type="checkbox"/> 3rd dose			<input type="checkbox"/> 4th dose		
<b>Haemophilus influenzae type b<sup>2</sup> (Hib)</b> <i>Protects against Haemophilus influenzae type b</i>			<input type="checkbox"/> 1st dose	<input type="checkbox"/> 2nd dose	<input type="checkbox"/> See footnote <sup>2</sup>		<input type="checkbox"/> 3rd or 4th dose, see footnote <sup>2</sup>			
<b>Pneumococcal (PCV13)</b> <i>Protects against pneumonia</i>			<input type="checkbox"/> 1st dose	<input type="checkbox"/> 2nd dose	<input type="checkbox"/> 3rd dose		<input type="checkbox"/> 4th dose			

You don't need to start over if your child misses a shot. Simply go back to your child's doctor for the next shot. Talk with your child's doctor if you have any questions about vaccines.

## RECOMMENDED IMMUNIZATIONS *continued...*

	Birth	1 month	2 months	4 months	6 months	9 months	12 months	15 months	18 months	23 months
<b>Inactivated poliovirus (IPV)</b> <i>Protects against polio</i>			<input type="checkbox"/> 1st dose	<input type="checkbox"/> 2nd dose	<input type="checkbox"/> 3rd dose					
<b>Influenza (Flu)<sup>3</sup></b> <i>Protects against the flu</i>					<input type="checkbox"/> Once every flu season <sup>3</sup>					
<b>Measles, mumps and rubella (MMR)</b> <i>Protects against measles, mumps and rubella</i>							<input type="checkbox"/> 1st dose			
<b>Varicella (VAR)</b> <i>Protects against chickenpox</i>							<input type="checkbox"/> 1st dose			
<b>Hepatitis A<sup>4</sup> (HepA)</b> <i>Protects against hepatitis A</i>							<input type="checkbox"/> Two-dose series, see footnote <sup>4</sup>			

## GROWTH *At each well-child visit, enter your child's measurements to keep track of their progress.*

Visit date										
Weight/Percentile	/	/	/	/	/	/	/	/	/	/
Length/Percentile	/	/	/	/	/	/	/	/	/	/
Head circumference										

 **FOR A CHECKLIST OF THINGS TO TALK ABOUT WITH YOUR CHILD'S DOCTOR AT THE NEXT APPOINTMENT, GO TO [HAP.ORG/DOCTORVISIT](https://www.hap.org/doctorvisit).**

<sup>1</sup> If Rotarix<sup>®</sup> is used, administer a two-dose series at ages 2 and 4 months. If RotaTeq<sup>®</sup> is used, administer a three-dose series at ages 2, 4 and 6 months. If any dose in the series was RotaTeq or the vaccine product is unknown for any dose, a total of three doses of the Rotavirus vaccine should be administered.

<sup>2</sup> Administer a two- or three-dose Hib vaccine primary series and booster dose (dose three or four depending on vaccine used in primary series).

<sup>3</sup> Two doses given at least four weeks apart are recommended for children starting at ages 6 months through 8 years who are getting a flu vaccine for the first time.

<sup>4</sup> Two doses of HepA vaccine are needed for lasting protection. The first dose should be given between 12 months and 23 months of age. The second dose should be given six to 18 months later.